

# AP1 Physics

Name \_\_\_\_\_

## Summer Assignment

Date \_\_\_\_\_ Per \_\_\_\_\_

### Fall 2022 – Spring 2023

**We have read the policies and expectations for AP Physics.**

**We understand and accept these policies.**

Student Signature: \_\_\_\_\_ Date \_\_\_\_\_

Parent / Guardian Name (print) \_\_\_\_\_

Parent / Guardian Signature: \_\_\_\_\_ Date \_\_\_\_\_

---

- I. As stated in the preceding “Course Policies and Expectations” sheet, all New Jersey Advanced Placement Classes are at a disadvantage. Many states and school districts follow the college schedule where classes begin in August and end in May. So the national exams are geared to these schools. You must take the exam one month early. This necessitates a very fast pace. This summer homework will allow us to start on the Physics subject matter immediately when school begins. This packet is a math review to brush up on valuable skills, and perhaps a means to assess whether you are correctly placed in Advanced Placement Physics.
- II. Physics, and AP Physics in particular, requires an exceptional proficiency in algebra, trigonometry, and geometry. In addition to the science concepts Physics often seems like a course in applied mathematics. The following assignment includes mathematical problems that are considered routine in AP Physics. This includes knowing several key metric system conversion factors and how to employ them. Another key area in Physics is understanding vectors.
- III. The attached pages contain a brief review, hints, and example problems. It is hoped that combined with your previous math knowledge this assignment is merely a review and a means to brush up before school begins in the fall. Please read the text and instructions throughout.

#### IV. **What is due the first day of school?**

A. **Signed Class Policies and Expectations Sheet**

1. **Read the sheet (the previous two pages)**
2. **Complete the section at the top of this form and obtain appropriate signatures.**

B. **Problems 1 to 13 (pages 2 to 10) of this packet (Row 1 of the AP Physics Assignment Sheet).**

C. **Row 2 of the AP Physics Assignment Sheet.**

**Check out the Physics Text.**

- **Read Chapter 2, sections 2.1 to 2.6.**
- **At the back of each chapter is a set of questions followed by a set of problems. Do questions 1 to 6 (Q 1 - 6) and problems 1 to 4 and 7 to 9 (P 1-4, 7-9) at the back of chapter 2.**

D. **Signed Safety Contract**

1. **Read the attached safety contract (Attached at the end of the packet)**
2. **Complete all parts of the form and obtain appropriate signatures.**

V. ***There will be a test covering the math review on the second day of class.***

VI. What if I don't get all the problems or don't understand the instructions?

- A. **Simply do the best you can, but show some work / effort in order to receive credit.**
- B. **Come to class the first day with your questions, in order to resolve these issues prior to the test.**

- The following are ordinary physics problems. Place the answer in scientific notation when appropriate and simplify the units (Scientific notation is used when it takes less time to write than the ordinary number does. As an example 200 is easier to write than  $2.00 \times 10^2$ , but  $2.00 \times 10^8$  is easier to write than 200,000,000).
- Your Ti-84 calculator is equipped to handle scientific notation in the following way:  $6.02 \times 10^{23}$  can be entered by putting in 6.02 2<sup>nd</sup> EE 23 This is the correct way to do all scientific notation. On the screen it will appear as 6.02E23. This places the whole number into parentheses, so it removes a step in the PEMDAS method.
- Units are the key to science, so do your best to simplify and find the final units for each calculation.

a.  $T_s = 2\pi \sqrt{\frac{4.5 \times 10^{-2} \text{ kg}}{2.0 \times 10^3 \text{ kg/s}^2}} =$  \_\_\_\_\_

b.  $K = \frac{1}{2} (1.99 \times 10^{-31} \text{ kg}) (2.8 \times 10^8 \text{ m/s})^2$  \_\_\_\_\_

c.  $F = \left( 9.0 \times 10^9 \frac{\text{N} \cdot \text{m}^2}{\text{C}^2} \right) \frac{(3.2 \times 10^{-9} \text{ C})(9.6 \times 10^{-9} \text{ C})}{(0.32 \text{ m})^2} =$  \_\_\_\_\_

d.  $\frac{1}{R_p} = \frac{1}{4.5 \times 10^2 \Omega} + \frac{1}{9.4 \times 10^2 \Omega}$   $R_p =$  \_\_\_\_\_

e.  $e = \frac{1.7 \times 10^3 \text{ J} - 3.3 \times 10^3 \text{ J}}{17 \times 10^3 \text{ J}}$  \_\_\_\_\_

f.  $1.33 \sin 25.0^\circ = 1.50 \sin \theta$   $\theta =$  \_\_\_\_\_

g.  $K_{\text{max}} = (6.63 \times 10^{-34} \text{ J} \cdot \text{s})(7.09 \times 10^{14} \text{ s}) - 2.17 \times 10^{-19} \text{ J} =$  \_\_\_\_\_

h.  $\gamma = \frac{1}{\sqrt{1 - \frac{2.25 \times 10^8 \text{ m/s}}{3.00 \times 10^8 \text{ m/s}}}} =$  \_\_\_\_\_

4. Often problems on the AP exam are done with variables only. Solve for the variable indicated. Don't let the different letters confuse you. Manipulate them algebraically as though they were numbers.

a.  $v_f = v_i + at$  ,  $a =$  \_\_\_\_\_

g.  $B = \frac{\mu_o I}{2\pi r}$  ,  $r =$  \_\_\_\_\_

b.  $K = \frac{1}{2} kx^2$  ,  $x =$  \_\_\_\_\_

h.  $x_m = \frac{m\lambda L}{d}$  ,  $d =$  \_\_\_\_\_

c.  $T_p = 2\pi \sqrt{\frac{\ell}{g}}$  ,  $g =$  \_\_\_\_\_

i.  $pV = nRT$  ,  $T =$  \_\_\_\_\_

d.  $F_g = G \frac{m_1 m_2}{r^2}$  ,  $r =$  \_\_\_\_\_

j.  $\sin \theta_c = \frac{n_1}{n_2}$  ,  $\theta_c =$  \_\_\_\_\_

e.  $mgh = \frac{1}{2} mv^2$  ,  $v =$  \_\_\_\_\_

k.  $qV = \frac{1}{2} mv^2$  ,  $v =$  \_\_\_\_\_

f.  $x = x_o + v_o t + \frac{1}{2} at^2$  ,  $t =$  \_\_\_\_\_

l.  $\frac{1}{f} = \frac{1}{s_o} + \frac{1}{s_i}$  ,  $s_i =$  \_\_\_\_\_

5. Science uses the **KMS** system (**SI**: System Internationale). **KMS** stands for kilogram, meter, second. These are the units of choice of physics. The equations in physics depend on unit agreement. So you must convert to **KMS** in most problems to arrive at the correct answer.

kilometers (*km*) to meters (*m*) and meters to kilometers  
 centimeters (*cm*) to meters (*m*) and meters to centimeters  
 millimeters (*mm*) to meters (*m*) and meters to millimeters  
 nanometers (*nm*) to meters (*m*) and meters to nanometers  
 micrometers ( $\mu m$ ) to meters (*m*)

gram (*g*) to kilogram (*kg*)  
 Celsius ( $^{\circ}C$ ) to Kelvin (*K*)  
 atmospheres (*atm*) to Pascals (*Pa*)  
 liters (*L*) to cubic meters ( $m^3$ )

Other conversions will be taught as they become necessary.

What if you don't know the conversion factors? Colleges want students who can find their own information (so do employers). Hint: Try a good dictionary and look under "measure" or "measurement". Or the Internet? Enjoy.

- |                                  |                     |                                   |                   |
|----------------------------------|---------------------|-----------------------------------|-------------------|
| a. 4008 <i>g</i>                 | = _____ <i>kg</i>   | h. 25.0 $\mu m$                   | = _____ <i>m</i>  |
| b. 1.2 <i>km</i>                 | = _____ <i>m</i>    | i. 2.65 <i>mm</i>                 | = _____ <i>m</i>  |
| c. 823 <i>nm</i>                 | = _____ <i>m</i>    | j. 8.23 <i>m</i>                  | = _____ <i>km</i> |
| d. 298 <i>K</i>                  | = _____ $^{\circ}C$ | k. 5.4 <i>L</i>                   | = _____ $m^3$     |
| e. 0.77 <i>m</i>                 | = _____ <i>cm</i>   | l. 40.0 <i>cm</i>                 | = _____ <i>m</i>  |
| f. $8.8 \times 10^{-8}$ <i>m</i> | = _____ <i>mm</i>   | m. $6.23 \times 10^{-7}$ <i>m</i> | = _____ <i>nm</i> |
| g. 1.2 <i>atm</i>                | = _____ <i>Pa</i>   | n. $1.5 \times 10^{11}$ <i>m</i>  | = _____ <i>km</i> |

6. Solve the following geometric problems.


- a. Line **B** touches the circle at a single point. Line **A** extends through the center of the circle.

- i. What is line **B** in reference to the circle?

\_\_\_\_\_


- ii. How large is the angle between lines **A** and **B**?

\_\_\_\_\_




- b. What is angle **C**?

\_\_\_\_\_




- c. What is angle  $\theta$ ?

\_\_\_\_\_



- d. How large is  $\theta$ ?

\_\_\_\_\_




- e. The radius of a circle is 5.5 cm,  
i. What is the circumference in meters?

\_\_\_\_\_


ii. What is its area in square meters?

- f. What is the area under the curve at the right?

\_\_\_\_\_



7. Using the generic triangle to the right, Right Triangle Trigonometry and Pythagorean Theorem solve the following. **Your calculator must be in degree mode.**



- a.  $\theta = 55^\circ$  and  $c = 32$  m, solve for  $a$  and  $b$ .

- d.  $a = 250$  m  
and  $b = 180$  m, solve for  $\theta$  and  $c$ .

- b.  $\theta = 45^\circ$  and  $a = 15$  m/s, solve for  $b$  and  $c$ .

- e.  $a = 25$  cm and  $c = 32$  cm, solve for  $b$  and  $\theta$ .

- c.  $b = 17.8$  m and  $\theta = 65^\circ$ , solve for  $a$  and  $c$ .

- f.  $b = 104$  cm and  $c = 65$  cm, solve for  $a$  and  $\theta$ .

\_\_\_\_\_

\_\_\_\_\_

## Vectors

Most of the quantities in physics are vectors. **This makes proficiency in vectors extremely important.**

**Magnitude:** Size or extend. The numerical value.

**Direction:** Alignment or orientation of any position with respect to any other position.

**Scalars:** A physical quantity described by a single number and units. A quantity described by **magnitude only.**

Examples: time, mass, and temperature

**Vector:** A physical quantity with **both a magnitude and a direction.** A directional quantity.

Examples: velocity, acceleration, force

Notation:  $\vec{A}$  or  $\vec{A}$

Length of the arrow is proportional to the vectors magnitude.

Direction the arrow points is the direction of the vector.

## Negative Vectors

Negative vectors have the same magnitude as their positive counterpart. They are just pointing in the opposite direction.



## Vector Addition and subtraction

Think of it as vector addition only. The result of adding vectors is called the resultant.  $\vec{R}$

$$\vec{A} + \vec{B} = \vec{R} \quad \vec{A} \text{ (arrow)} + \vec{B} \text{ (arrow)} = \vec{R} \text{ (arrow)}$$

So if  $A$  has a magnitude of 3 and  $B$  has a magnitude of 2, then  $R$  has a magnitude of  $3+2=5$ .

When you need to subtract one vector from another think of the one being subtracted as being a negative vector. Then add them.

$\vec{A} - \vec{B}$  is really

$$\vec{A} + \vec{B} = \vec{R}$$


A negative vector has the same length as its positive counterpart, but its direction is reversed. So if  $A$  has a magnitude of 3 and  $B$  has a magnitude of 2, then  $R$  has a magnitude of  $3+(-2)=1$ .

**This is very important.** In physics a negative number does not always mean a smaller number. Mathematically  $-2$  is smaller than  $+2$ , but in physics these numbers have the same magnitude (size), they just point in different directions ( $180^\circ$  apart).


There are two methods of adding vectors

**Parallelogram**

$A + B$




$A - B$




**Tip to Tail**

$A + B$




$A - B$



It is readily apparent that both methods arrive at the exact same solution since either method is essentially a parallelogram. It is useful to understand both systems. In some problems one method is advantageous, while in other problems the alternative method is superior.

8. Draw the resultant vector using the parallelogram method of vector addition.


Example




b.




d.




a.



c.




e.




9. Draw the resultant vector using the tip to tail method of vector addition. Label the resultant as vector  $R$


Example 1:  $A + B$




Example 2:  $A - B$




a.  $X + Y$




d.  $C - D$




b.  $T - S$




e.  $A + B + C$






c.  $P + V$



f.  $A - B - C$



**Direction:** What does positive or negative direction mean? How is it referenced? The answer is the coordinate axis system. **In physics a coordinate axis system is used to give a problem a frame of reference.** Positive direction is a vector moving in the positive  $x$  or positive  $y$  direction, while a negative vector moves in the negative  $x$  or negative  $y$  direction (This also applies to the  $z$  direction, which will be used sparingly in this course).





What about vectors that don't fall on the axis? You must specify their direction using degrees measured from East.


### Component Vectors

A resultant vector is a vector resulting from the sum of two or more other vectors. Mathematically the resultant has the same magnitude and direction as the total of the vectors that compose the resultant. Could a vector be described by two or more other vectors? Would they have the same total result?

This is the reverse of finding the resultant. You are given the resultant and must find the component vectors on the coordinate axis that describe the resultant.




Or




Any vector can be described by an  $x$  axis vector and a  $y$  axis vector which summed together mean the exact same thing. The advantage is you can then use plus and minus signs for direction instead of the angle.

10. For the following vectors draw the component vectors along the  $x$  and  $y$  axis.


a.




c.



b.





d.



Obviously the quadrant that a vector is in determines the sign of the  $x$  and  $y$  component vectors.

### Trigonometry and Vectors


Given a vector, you can now draw the  $x$  and  $y$  component vectors. The sum of vectors  $x$  and  $y$  describe the vector exactly. Again, any math done with the component vectors will be as valid as with the original vector. The advantage is that math on the  $x$  and/or  $y$  axis is greatly simplified since direction can be specified with plus and minus signs instead of degrees. But, how do you mathematically find the length of the component vectors? Use trigonometry.



$$\begin{aligned} \cos \theta &= \frac{adj}{hyp} & \sin \theta &= \frac{opp}{hyp} \\ adj &= hyp \cos \theta & opp &= hyp \sin \theta \\ x &= hyp \cos \theta & y &= hyp \sin \theta \\ x &= 10 \cos 40^\circ & y &= 10 \sin 40^\circ \\ x &= 7.66 & y &= 6.43 \end{aligned}$$

11. Solve the following problems. You will be converting from a polar vector, where direction is specified in **degrees measured counterclockwise from east**, to component vectors along the  $x$  and  $y$  axis. Remember the plus and minus signs on you answers. They correspond with the quadrant the original vector is in.  
 Hint: Draw the vector first to help you see the quadrant. Anticipate the sign on the  $x$  and  $y$  vectors. Do not bother to change the angle to less than  $90^\circ$ . Using the number given will result in the correct + and - signs. The first number will be the magnitude (length of the vector) and the second the degrees from east.  
**Your calculator must be in degree mode.**

Example: 250 at  $235^\circ$



$$x = \text{hyp} \cos \theta$$

$$x = 250 \cos 235^\circ$$

$$x = -143$$

$$y = \text{hyp} \sin \theta$$

$$y = 250 \sin 235^\circ$$

$$y = -205$$

Another and more useful way to do vectors is with North, East, South and West quadrants. This is more applicable to real life situations, such as air, sea and land travel. It is also how buildings and structures are oriented in Civil Engineering and Architecture.

If the top of the page is considered North, then the above example could have been listed as  $55^\circ$  South of West. When we review this in class, we will most likely do it this way, as it is the most sensible. X is considered East, -X is West, Y is North, -Y is South. Very logical.

For the following, please show the x and y components of each:

a. 89 at  $150^\circ$

b. 6.50 at  $345^\circ$

c. 0.00556, at  $60^\circ$  South of West

d.  $7.5 \times 10^4$ ,  $30^\circ$  North of East




e. 12 at 265°

f. 990 at 320°

12. 8653 at 225° Given two component vectors solve for the resultant vector. This is the opposite of number 11 above. Use Pythagorean Theorem to find the hypotenuse, then use inverse (arc) tangent to solve for the angle.

Example:  $x = 20$ ,  $y = -15$



$$R^2 = x^2 + y^2 \quad \tan \theta = \frac{opp}{adj}$$

$$R = \sqrt{x^2 + y^2} \quad \theta = \tan^{-1} \frac{opp}{adj}$$

$$R = \sqrt{20^2 + 15^2}$$

$$\theta = \tan^{-1} \frac{y}{x}$$

$$R = 25$$

$$\theta = \tan^{-1} \left( \frac{-15}{20} \right) = -36.9^\circ$$

$$360^\circ - 36.9^\circ = 323.1^\circ$$

Or this could be given as 25 units, 36.9°, South of East

a.  $x = 600$ ,  $y = 400$

c.  $x = -32$ ,  $y = 16$

b.  $x = -0.75$ ,  $y = -1.25$

d.  $x = 0.0065$ ,  $y = -0.0090$

e.  $x = 20,000, y = 14,000$

f.  $x = 325, y = 998$

## How are vectors used in Physics?

They are used everywhere!

### Speed

Speed is a scalar. It only has magnitude (numerical value).

$v_s = 10 \text{ m/s}$  means that an object is going 10 meters every second. But, we do not know where it is going.

### Velocity

Velocity is a vector. It is composed of both magnitude and direction. Speed is a part (numerical value) of velocity.

$v = 10 \text{ m/s}$  north, or  $v = 10 \text{ m/s}$  in the  $+x$  direction, etc.

There are three types of speed and three types of velocity

**Instantaneous speed / velocity:** The speed or velocity at an instant in time. You look down at your speedometer and it says  $20 \text{ m/s}$ . You are traveling at  $20 \text{ m/s}$  at that instant. Your speed or velocity could be changing, but at that moment it is  $20 \text{ m/s}$ .

**Average speed / velocity:** If you take a trip you might go slow part of the way and fast at other times. If you take the total distance traveled divided by the time traveled you get the average speed over the whole trip. If you looked at your speedometer from time to time you would have recorded a variety of instantaneous speeds. You could go  $0 \text{ m/s}$  in a gas station, or at a light. You could go  $30 \text{ m/s}$  on the highway, and only go  $10 \text{ m/s}$  on surface streets. But, while there are many instantaneous speeds there is only one average speed for the whole trip.

**Constant speed / velocity:** If you have cruise control you might travel the whole time at one constant speed. If this is the case then your average speed will equal this constant speed.

#### A trick question

Will an object traveling at a constant speed of  $10 \text{ m/s}$  also always have constant velocity?

Not always. If the object is turning around a curve or moving in a circle it can have a constant speed of  $10 \text{ m/s}$ , but since it is turning, its direction is changing. And if direction is changing then velocity must change, since velocity is made up of speed and direction.

**Constant velocity must have both constant magnitude and constant direction.**

### Rate

Speed and velocity are rates. A rate is a way to quantify anything that takes place during a time interval. Rates are easily recognized. They always have time in the denominator.

$10 \text{ m/s}$                        $10 \text{ meters / second}$

## The very first Physics Equation

Velocity and Speed both share the same equation. Remember speed is the numerical (magnitude) part of velocity. Velocity only differs from speed in that it specifies a direction.

$$v = \frac{x}{t} \quad v \text{ stands for velocity} \quad x \text{ stands for displacement} \quad t \text{ stands for time}$$

**Displacement** is a vector for distance traveled in a straight line. It goes with velocity. Distance is a scalar and goes with speed. Displacement is measured from the origin. It is a value of how far away from the origin you are at the end of the problem. The direction of a displacement is the shortest straight line from the location at the beginning of the problem to the location at the end of the problem.

How do distance and displacement differ? Suppose you walk 20 meters down the  $+x$  axis and turn around and walk 10 meters down the  $-x$  axis.

The distance traveled does not depend on direction since it is a scalar, so you walked  $20 + 10 = 30$  meter.

Displacement only cares about your distance from the origin at the end of the problem.  $+20 - 10 = 10$  meter.

13. Attempt to solve the following problems. Take heed of the following.

**Always use the *KMS* system: Units must be in kilograms, meters, seconds.**

**On the all tests, including the AP exam you must:**

**1. List the original equation used.**

**2. Show correct substitution.**

**3. Arrive at the correct answer with correct units.**

Distance and displacement are measured in meters (m)

Speed and velocity are measured in meters per second (m/s)

Time is measured in seconds (s)

Example: A car travels 1000 meters in 10 seconds. What is its velocity?

$$v = \frac{x}{t} \qquad v = \frac{1000m}{10s} \qquad v = 100m/s$$

- a. A car travels 35 km west and 75 km east. What distance did it travel?
  
  
  
  
  
  
  
  
  
  
- b. A car travels 35 km west and 75 km east. What is its displacement?
  
  
  
  
  
  
  
  
  
  
- c. A car travels 35 km west, 90 km north. What distance did it travel?
  
  
  
  
  
  
  
  
  
  
- d. A car travels 35 km west, 90 km north. What is its displacement?
  
  
  
  
  
  
  
  
  
  
- e. A bicyclist pedals at 10 m/s in 20 s. What distance was traveled?
  
  
  
  
  
  
  
  
  
  
- f. An airplane flies 250.0 km at 300 m/s. How long does this take?
  
  
  
  
  
  
  
  
  
  
- g. A skydiver falls 3 km in 15 s. How fast are they going?
  
  
  
  
  
  
  
  
  
  
- h. A car travels 35 km west, 90 km north in two hours. What is its average speed?
  
  
  
  
  
  
  
  
  
  
- i. A car travels 35 km west, 90 km north in two hours. What is its average velocity?

- j. You take a trip to the shore, and travel at 60 mph the whole way down. Then, on the way back, you hit traffic, and travel at 30 mph the whole way home. What was your average speed for the trip?